



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

yeast apparently has a surface tension of about 0.60. The article seems to clear away much of the haze that has surrounded the matter of the significance of surface tension in cell activity.—WILLIAM CROCKER.

**Underground organs of weeds.**—Conflicting statements by various authors have induced PAMMEL and FOGEL<sup>23</sup> to investigate the organs of vegetative reproduction of some of our most common weeds. The Canada thistle (*Cirsium arvense*), the horse nettle (*Solanum carolinense*), the milkweed (*Asclepias syriaca*), and the bindweed (*Convolvulus arvensis*), were all found to be propagated by horizontal roots bearing adventitious buds; while in the wild morning glory (*Convolvulus Sepium*) and the quack grass (*Agropyron repens*), the organs of vegetative multiplication are rootstocks. In some instances the roots and subterranean stems resembled each other so closely that only by microscopic examination could the difference be detected.—GEO. D. FULLER.

**Epidermis and light refraction.**—FRIMMEL<sup>24</sup> thinks he has shown that the lower papillate epidermis of the leaves of the yew gives a total refraction of the light passing through the leaf from above, thereby leading to the use of all light that enters the leaf. He relates this character to the ability of the tree to grow in shaded habitats. He believes the lower epidermis of a number of other conifers acts in the same way. He finds a similar contrivance in the spongy parenchyma of the cotyledon of the beech. The fact of total refraction in the yew seems entirely established; whether it is of biological significance or not is quite another question.—WILLIAM CROCKER.

**Arctic vegetation.**—Hare Island off the coast of West Greenland, an uninhabited island 66 square miles in area, has been visited several times by PORSILD,<sup>25</sup> who has found a flora consisting of 82 arctic and 30 subarctic species. The vegetation belongs to the fell-field formation, large areas quite devoid of plants, passing into a poorly developed heath with arctic meadows and bogs in the more sheltered situations. Dispersal is almost entirely through the agency of the wind over the surface of the snow and frozen seas. The sub-arctic species are regarded as relics of milder climate in post-glacial times.—GEO. D. FULLER.

**Pneumatophores.**—From an examination of the tissues of vertical apogeo-tropic branches of the roots of *Terminalia Arjuna*, a large tree of Central India,

<sup>23</sup> PAMMEL, L. H., and FOGEL, ESTELLE D., The underground organs of a few weeds. Proc. Iowa Acad. Sci. 16: pp. 7. pls. 5. 1909.

<sup>24</sup> FRIMMEL, FRANZ V., Die untere Kutikula des *Taxus*-Blattes ein Lichtreflektor Oester. Bot. Zeitsch. 61: 216-223. figs. 4. 1911.

<sup>25</sup> PORSILD, MORTEM P., The plant life of Hare Island off the coast of West Greenland. Særtryk af Meddelelser om Gronland 47: 252-274. figs. 10. Kobenhavn: Bianco Lunos. 1910.

ADAMSON<sup>26</sup> decides that they are developed "for purposes of aeration as shown by the great development of lacunar tissue." Both the horizontal and the vertical roots possess very loose cortical tissue with large lacunae, but most botanists would probably hesitate to pronounce upon the purpose of its development. The upright roots have well developed root caps, and possess no lenticels or other stem characters found in many pneumatophores.—GEO. D. FULLER.

**Seed of *Neuropteris*.**—In 1904 KIDSTON described the seed of *Neuropteris heterophylla*, which was said to be "as large as a hazelnut." Now the same investigator, associated with JONGMANS, has described<sup>27</sup> the seed of *N. obliqua* Brong., the specimens being in the Rijks Herbarium at Leyden. The seeds have the same general structure as those of *N. heterophylla*, but are about twice as large. This species of *Neuropteris* is also doubtless to be referred to the stem genus *Medullosa*.—J. M. C.

**Root parasites.**—MISS BENSON<sup>28</sup> has studied the structure of some haustoria on the roots of *Exocarpus* and *Thesium*, showing the nature of the penetration and connection with the roots of other plants. For a portion of the lignified elements of the haustoria the name "phloeotracheids" is suggested, and the investigator thinks they may act as a filter between the host and parasite, although the evidence that they have any such function does not seem to be at all convincing.—GEO. D. FULLER.

**Calcium salts and fungi.**—WEIR<sup>29</sup> concludes that soluble calcium salts are necessary to the complete development of higher fungi. *Coprinus plicatilis*, *C. papillatus*, *C. nivens*, and *C. ephemoides* showed little if any mycelial development, and no development of fruit heads or spores, when all the calcium present was in the form of the oxalate.—WILLIAM CROCKER.

**A bog in central Illinois.**—GATES<sup>30</sup> has instanced the meeting of northern and southern forms in a bog in central Illinois.—GEO. D. FULLER.

<sup>26</sup> ADAMSON, R. S., Note on the roots of *Terminalia Arjuna*. New Phytol. **9**: 150-156. figs. 3-7. 1910.

<sup>27</sup> KIDSTON, R., and JONGMANS, W. J., Sur la fructification de *Neuropteris obliqua* Bgt. Archiv. Néerl. Sci. III. B. **1**: 25, 26. pl. 1. 1911.

<sup>28</sup> BENSON, MARGARET, Root parasitism in *Exocarpus* (with comparative notes on the haustoria of *Thesium*). Ann. Botany **24**: 667-677. pl. 65. figs. 4. 1910.

<sup>29</sup> WEIR, JAMES R., Benötigt der Pilz *Coprinus* Kalksalze zu seinen physiologischen Funktionen. Flora **103**: 87-90. 1911.

<sup>30</sup> GATES, F. C., A bog in central Illinois. Torreya **11**: 205-211. 1911.